


The next ten years of core-collapse supernova gravitational-wave astronomy

Marek Szczepańczyk
Department of Physics,
University of Warsaw

<https://www.fuw.edu.pl/~mszczepanczyk/>

Slides:




Stellar Populations and their Explosions:
Bridging the Gap
Schloss Ringberg, Tegernsee, 13.11.2025

10 years of Gravitational-Wave Astronomy!

GW150914: discovery of gravitational waves on 14 Sep 2015


What's next?



Outline

- Preliminaries
 - Observing timelines
 - Detection distance
- Are we ready (as communities) for a nearby CCSN?
 - SN2025gw IGWN Symposium
 - Some lessons learned

Past and Future 10 years



- <https://observing.docs.ligo.org/plan/>
- O4: 24 months total, until 18 Nov 2025
- LIGO: up to 180 Mpc, currently 160 Mpc
- Virgo: around 55 Mpc
- KAGRA: reached 7.5 Mpc, observed briefly

- Also: GEO600, a few Mpc
- LIGO-India - under construction
- Possible an Australian detector

Binaries and binaries so far...


- Low-latency searches are crucial for astronomical observations, resources:
 - <https://gracedb.ligo.org>
 - <https://emfollow.docs.ligo.org/userguide/>
- Published results:
 - 21 papers and counting!
 - GWTC-4, O4a catalog ([2508.18082](#))
 - Search with SN 2023ixf, a core-collapse supernova at 6.7 Mpc ([2410.16565](#))

GW candidates so far (**3 per week**):

O4a: 128 GWs



O4b: 105 GW candidates

O4c: 66 GW candidates






Optically targeted searches

- While waiting for a Galactic CCSN, we can systematically constrain its engine with CCSNe at Mpc range -> **optically targeted searches**
- **O1-O2 search** (Abbott+19, [1908.03584](#)):
 - First observational constraints of a CCSN engine
- **O3 search** (Szczepanczyk+24, [2305.16146](#)):
 - We could not beat the previous upper limits
- **SN 2023ixf search** (Abac+24, [2410.16565](#), special O4 paper):
 - GW energy emission: constraints improved by an order of magnitude





When will we discover GWs? (realistically: Galactic CCSN)



2.5G detectors (post-O5, before 3G)

- Szczepańczyk&Zanolin 2022 *Gravitational Waves from a Core-Collapse Supernova: Perspectives with Detectors in the Late 2020s and Early 2030s*, [Galaxies 2022, 10\(3\), 70](#)
- Detectors: LIGO Hanford, LIGO Livingston, LIGO India, Virgo, KAGRA, NEMO/Australia (?)
- Factor of 2 with respect to 2G detectors (conservative)
- Improved sky localization



Are we ready, as
communities, for a CCSN
in Milky Way (or nearby)?

SN2025gw: IGWN Symposium on CCSNe Bridging the Gap

- IGWN - International Gravitational-Wave Network, to replace LVK
- Symposium webpage: <https://indico2.fuw.edu.pl/event/17/overview>
- CCSNe are the most challenging astronomical events to model
 - Theory, GW, neutrino, EM, next 10 years
- Slack channel: please ask me to add you
- Youtube: [playlist](#)
- Large attendance of the Symposium: **Galactic CCSN - the next big thing?**



SN2025gw: IGWN Symposium on CCSNe

- Classical and Quantum Gravity focus issue proceedings and White Paper (in progress): <https://iopscience.iop.org/collections/cqg-250513-841>
- A White Paper will summarize the state of the art of the fields and provide recommendations
- **Everybody can contribute**, regardless of attending the Symposium:
 - Regular articles
 - Proceedings papers
- Deadline (likely soft): 31 Dec 2025

Classical and Quantum Gravity

Focus on Core Collapse Supernova Gravitational Wave Astronomy and Astrophysics: Past, Present, and Future

Guest Editors

Marek Szczepańczyk, *University of Warsaw, Poland*

Marco Cavaglia, *Missouri University of Science and Technology, United States*

Anthony Mezzacappa, *University of Tennessee Knoxville, United States*

Jade Powell, *Swinburne University of Technology, Australia*

Scope

The LIGO, Virgo, and KAGRA (LVK) observatories are designed to detect gravitational waves (GWs) from a wide variety of sources, including compact binary mergers, core-collapse supernovae, and isolated pulsars, among others. Only the

SN2025gw: IGWN Symposium on CCSNe


Some lessons learned (work in progress on a White Paper):

- **A multimessenger (GW+nu+EM) detection program** for CCSNe is underway. Detailed discussion/decisions needed on the conditions for declaring a detection with a CCSN candidate (statistical significance, morphological constraints).
- **A multimessenger (GW+nu+EM) parameter estimation program** for CCSNe is underway. More areas can be expanded further like multi feature.
- **The interaction between EM, nu and GW communities could be improved**, for example what will we do explicitly if we have a Galactic CCSNe?
- While there is a consensus on predicted GWs, it is still not known what waveform amplitudes we may expect.
- Many uncertainties (**progenitor structure...**) and hidden assumptions (**transport treatment...**) are baked into the multi-physics supernova problem.
- Important to consider how supernova modelling needs to **evolve technically and organisationally** to make further progress and aid gravitational wave and multi-messenger astronomy.


FAQ: can LVK run continuously?

Short answer: (we wish it too but) no..., because

- „**You cannot eat cookie and have cookie**”: you either make observations at a given sensitivity or do breaks between runs to improve the sensitivity (we do not want to run at O1 sensitivity for 20 years)
- **Detector upgrades**: replacing hardware, testing, calibrating
- **Interruptions during a run**: weather, maintenance, cleaning, equipment failures/malfunctioning, replenishing technical gases, locking etc.
- **Coincidence duty factor**: synchronizing detectors to get double and triple coincidence
- **Single detector data**: hard case for GW bursts (e.g. Cavaglia+20)



Papers on GWs from CCSN



CCSN GW Literature, by Ewald Mueller:


https://wwwmpa.mpa-garching.mpg.de/rel_hydro/GWlit_catalog.shtml

Low-latency Physical Inference work in progress

Approach:

Implement in low-latency


Existing methods



Summary


- Next 10 years of GW detectors
 - Pushing facilities and prepare for ET/CE
- Are we ready?
 - *SN2025gw: IGWN Symposium on CCSNe*, first such a meeting of multimessenge community, likely first meeting of a series
 - White paper and proceedings in progress. Everybody is welcome to contribute!

“Welcome SN 202X! Long-awaited for 2025-2026”
Fukuoka Temple, 2019.10.23




Extras

CCSN - parameter estimation



GW detection challenges


More details: Szczepańczyk+21 ([2104.06462](#))




- Large time-frequency area, in contrast to GWs from BBHs
- Detectable signal-to-noise ratio of 20-30

Model-independent searches

- **Coherent WaveBurst** (cWB, Klimenko+16) is a software designed to detect a wide range of burst transients without prior knowledge of the signal morphology
- cWB uses minimal assumptions, for example growing frequency over time in case of binaries
- **Complementing template-based searches**
- cWB has detected:
 - **GW150914** - the very first GW (PRL 116, 061102)
 - **GW190521** - an intermediate mass binary black hole, 150 Msun (PRL 125, 101102)
 - **GW231123** - ~200 Msun BH ([2507.08219](https://arxiv.org/abs/2507.08219))
 - It regularly detects GWs together with template-based searches
- The cWB contributes results to several LVK papers during each observing run.




<https://gwburst.gitlab.io/>



Model-independent searches classification


Compact binary searches (minimally modeled)

Binary black holes
Binary neutron stars
Black hole - neutron star



e.g. Mishra+23 ([2201.01495](#))


Binaries with eccentric orbits
Intermediate-mass black holes
Hyperbolic encounters
Extreme mass-ratio



e.g. MS+21 ([2009.11336](#))

Generic searches (unmodeled)

Core-collapse supernovae
Pulsar glitches
Cosmic strings
Unknown



e.g. MS+24 ([2305.16146](#))


Low-latency searches



Public alerts for
multi-messenger observations:
electromagnetic, cosmic rays,
and neutrino

e.g. Chaudhary+24 ([2308.04545](#))

Searches for new phenomena



Higher harmonics
GW cross-polarization
Deviations from GR

e.g. Vedovato+22 ([2108.13384](#))

IGWN Symposium on CCSNe

(IGWN - International GW Network)

Some lessons learned (work in progress on a White Paper):

- 2D simulations can be reliable for computing the frequency of the high-frequency feature, 3D are needed for the correct amplitudes.
- There is more effort in computing the GW emission for rapid rotation CCSNe. This is better than a few years ago, but we need to have quantitative comparisons
- The Core Bounce Phase of rapid rotation is templetale (still a few percent uncertainty in the amplitude).
- Some debate on the relationship of the SASI frequency in the GW and neutrino channel (f_{gw} twice f_{nu} is the leading expectation).
- Which mode is related to the high-frequency feature depends on the progenitor and simulations. Not clear yet how well we can distinguish between f and g modes from GW (modal analysis can be used to interpret the frequency evolution if the mode is established, but does not predict amplitudes)
- Usage of EM information is still in early phase. We currently use light curved for estimating the time of the shock breakout but many other areas can be used/expanded for MMA with CCSNe (spectra, progenitor mass estimation, nucleosynthesis).
- Some people are starting to work on “less unmodelled” configurations for CCSNe GW searches.

Neutrino data

Simulation:


200 GB

Reduced Data:

5 GB

Marek needs:

2 Bytes



(Observational)

time of the collapse

